



Contents lists available at ScienceDirect

The Journal of Arthroplasty

journal homepage: www.arthroplastyjournal.org

Health Policy & Economics

Predictors and Cost of Readmission in Total Knee Arthroplasty



Kenneth L. Urish, MD, PhD ^{a,*}, Yongmei Qin, MD, MS ^b, Benjamin Y. Li, BS ^b,
 Tudor Borza, MD, MS ^b, Michael Sessine, BS ^b, Peter Kirk, MS ^b,
 Brent K. Hollenbeck, MD, MS ^b, Jonathan E. Helm, PhD ^c, Mariel S. Lavieri, PhD ^d,
 Ted A. Skolarus, MD, MPH ^e, Bruce L. Jacobs, MD, MPH ^f

^a Arthritis and Arthroplasty Design Group, The Bone and Joint Center, Magee Womens Hospital of the University of Pittsburgh Medical Center; Department of Orthopaedic Surgery, Department of Bioengineering, and Clinical and Translational Science Institute, University of Pittsburgh; Department of Biomedical Engineering, Carnegie Mellon University, Pittsburgh, Pennsylvania

^b Division of Oncology, Dow Division for Urologic Health Service Research, Department of Urology, University of Michigan, Ann Arbor, Michigan

^c Health Services Research & Development, Center for Clinical Management Research, VA Ann Arbor Healthcare System, Ann Arbor, Michigan

^d Department of Industrial and Operations Engineering, University of Michigan, Ann Arbor, Michigan

^e Department of Urology, Division of Oncology, Dow Division for Urologic Health Service Research, Health Services Research & Development, Center for Clinical Management Research, VA Ann Arbor Healthcare System, University of Michigan, Ann Arbor, Michigan

^f Department of Urology, University of Pittsburgh, Pittsburgh, Pennsylvania

ARTICLE INFO

Article history:

Received 23 February 2018

Received in revised form

25 March 2018

Accepted 2 April 2018

Available online 17 April 2018

Keywords:

total knee arthroplasty

readmission

Comprehensive Care for Joint Replacement

Nationwide Readmission Database

readmission cost

ABSTRACT

Background: The Comprehensive Care for Joint Replacement bundle was created to decrease total knee arthroplasty (TKA) cost. To help accomplish this, there is a focus on reducing TKA readmissions. However, there is a lack of national representative sample of all-payer hospital admissions to direct strategy, identify risk factors for readmission, and understand actual readmission cost.

Methods: We used the Nationwide Readmission Database to examine national readmission rates, predictors of readmission, and associated readmission costs for elective TKA procedures. We fit a multivariable logistic regression model to examine factors associated with readmission. Then, we determined mean readmission costs and calculated the readmission cost when distributed across the entire TKA population.

Results: We identified 224,465 patients having TKA across all states participating in the Nationwide Readmission Database. The mean unadjusted 30-day TKA readmission rate was 4%. The greatest predictors of readmission were congestive heart failure (odds ratio [OR] 2.51, 95% confidence interval [CI] 2.62–2.80), renal disease (OR 2.19, 95% CI 2.03–2.37), and length of stay greater than 4 days (OR 2.4, 95% CI 2.25–2.61). The overall median cost for each readmission was \$6753 ± 175. Extrapolating the readmission cost for the entire TKA population resulted in the readmission cost being 2% of the overall 30-day procedure cost.

Conclusions: A major focus of the Comprehensive Care for Joint Replacement bundle is improving cost and quality by limiting readmission rates. TKA readmissions are low and comprise a small percentage of total TKA cost, suggesting that they may not be the optimal measure of quality care or a significant driver of overall cost.

© 2018 Elsevier Inc. All rights reserved.

One or more of the authors of this paper have disclosed potential or pertinent conflicts of interest, which may include receipt of payment, either direct or indirect, institutional support, or association with an entity in the biomedical field which may be perceived to have potential conflict of interest with this work. For full disclosure statements refer to <https://doi.org/10.1016/j.arth.2018.04.008>.

Funding: Dr. Kenneth Urish is supported in part by the National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS K08AR071494), the National Center for Advancing Translational Science (NCATS KL2TR0001856), the Orthopaedic Research and Education Foundation, and the Musculoskeletal Tissue Foundation. Dr. Bruce Jacobs is supported in part by the National Institutes of Health Institutional KL2 award (KL2TR000146-08), the GEMSTAR award (R03AG048091), and the Jahnigen Career Development Award. Dr. Tudor Borza is supported by a National Cancer Institute Training Grant (T32-CA180984). Dr. Mariel Lavieri is

supported by the National Science Foundation (CMMI-1552545). Dr. Brent Hollenbeck is supported in part by Research Scholar Grant RSGI-13-323-01-CPHPS from the American Cancer Society and by NIH/NCI grant R01 CA168691, National Institute of Aging, Agency for Healthcare Research and Quality (R01 HS18726). Dr. Ted Skolarus is supported by a VA Health Services Research & Development Career Development Award-2 (CDA 12-171).

Dr. Brent Hollenbeck is also an Associate Editor of Urology.

* Reprint requests: Kenneth L. Urish, MD, PhD, Department of Orthopaedic Surgery, Department of Bioengineering, Department of Biomedical Engineering, Arthritis and Arthroplasty Design Group, The Bone and Joint Center, Magee Womens Hospital of the University of Pittsburgh Medical Center; Clinical and Translational Science Institute, University of Pittsburgh; Carnegie Mellon University, 300 Halket Street, Suite 1601, Pittsburgh, PA 15232.

<https://doi.org/10.1016/j.arth.2018.04.008>

0883-5403/© 2018 Elsevier Inc. All rights reserved.

As one of the largest procedural expenditures for Medicare, total knee arthroplasty (TKA) has been under intense scrutiny to decrease costs [1]. The Affordable Care Act of 2012 began a long anticipated shift in Medicare reimbursement policy from a fee-for-service model to one linked to quality of care [2]. As part of this effort, Medicare launched the Comprehensive Care for Joint Replacement (CJR) model to bundle all the hospital, professional, and 90-day postdischarge services into a single payment for TKA [3]. Currently, this bundled care pertains only to selected regions, but it is anticipated that it will soon expand across the United States. Following the pattern of Medicare, it is expected that private insurers will adopt this strategy. The logic behind bundled care involves tying all parties of the episode of care together to align their interests [4]. Hospitals in charge of the overall episode have already improved inpatient efficiency [5]. Understanding excess costs of postdischarge care has become essential, and reducing readmissions rates is an obvious and assumed initial target.

The readmission rates of TKA are low between 2% and 6%, and there are a large number of predictors of readmission [6]. Large databases that are able to track postdischarge readmissions and charge data are scarce, and these databases lack a national representative sample of all-payer hospital admissions across a wide geographic region. This is a disadvantage of using the National Inpatient Sample or Medicare databases, 2 large databases used to study TKA readmissions for health policy research. The Nationwide Readmissions Database (NRD) was developed to address these gaps. The NRD is the largest database of all-payer hospital admissions that can follow readmissions across different hospitals and accounts for 49% of all hospitalizations in the United States [7].

For these reasons, we used the NRD to examine national readmission rates, predictors of readmission, and associated readmission costs for elective TKA procedures. The objective was to quantify 30-day NRD readmission rates for TKA, identify predictors of readmission, and quantify readmission cost as a function of overall care for the population. We hypothesized that TKA readmission rates and costs associated with readmission aggregated over the entire population would be low. Findings from this study will inform policy makers about patterns of TKA readmissions and the actual contribution of readmission rate cost across all payers and geographic regions in the United States.

Methods

Data Source and Study Population

We utilized the Healthcare Cost and Utilization Project's NRD to identify adult men and women (18 years or older) who underwent TKA in 2014. The NRD was developed to provide a nationally representative data set to examine hospital readmissions [7]. Sampled from the State Inpatient Database, the NRD is the largest database of all-payer hospital admissions that contains patient linkages to follow readmissions across different hospitals [7]. Initial data were released for 2014 and first became available in 2016 [7].

We chose to study TKA because it is one of the conditions included in the recently implemented CJR bundle payment model wherein Medicare pays a hospital a single payment to cover the costs of hospital, professional, and postacute care services that occur within 90 days of the procedure [3]. We identified TKA using designated International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) codes. As there may be more than one procedure code for each patient, we included all patients with the ICD-9 procedure code 8154 and excluded any patients with the ICD-9 procedure code 8151, 8006, 8155, 0080, 0081, 0082, 0083, 0084, 27486, 27487, 274888 and diagnosis codes 73310, 73314, 73315, 73316, 808, 820, 821, 827, 828. Using these criteria, our study

consisted of 224,465 patients. These codes were used in an attempt to include elective TKA and exclude TKA performed for trauma or neoplasm.

Outcomes

The objective of this study was to assess readmission rates among patients undergoing TKA. First, we defined a readmission as a hospital admission that occurred within 30 days of the index surgery. We used a 30-day time period to be consistent with the readmission definition used by the Hospital Readmissions Reduction Program [8].

Statistical Analysis

We first compared patient demographics as well as clinical, payer, and hospital characteristics among TKA patients readmitted and not readmitted. We then examined readmitted patients stratified by payer type (ie, Medicare or non-Medicare). Non-Medicare payers included Medicaid, private insurance, uninsured, and other insurance types. Comorbidity was measured using the adaptation of the Charlson index by Deyo et al [9] in which ICD-9 diagnosis and procedure codes were used to identify the presence or absence of 16 comorbid conditions. Rural areas included micropolitan areas. A teaching hospital was defined as one that was a member of the Council of Teaching Hospitals, had an Accreditation Council for Graduate Medical Education–approved residency program, or had a ratio of full-time equivalent interns and residents to beds of 0.25 or higher [10]. Parametric continuous variables were compared using Student *t* tests, nonparametric continuous variables were compared using the Wilcoxon rank-sum test, and categorical variables were compared using chi-square tests.

Next, we fit univariable and multivariable logistic regression models to examine factors associated with readmission. For the multivariable model, covariates included age, sex, median household income, comorbidities, index admission length of stay, primary payer (ie, Medicare or non-Medicare), hospital type (ie, teaching or nonteaching), urban vs rural location, and hospital size. The effect of each covariate was adjusted for all other covariates in the model.

Finally, we focused on readmission costs. Specifically, we calculated readmission costs using an established method [11,12] based on total readmission charges and hospital-specific cost-to-charge ratios developed by the Healthcare Cost and Utilization Project [7]. The cost-to-charge ratio provides a way to estimate the cost of hospital services, as opposed to the charges put forth by hospitals. We determined mean readmission costs and then calculated what the readmission cost would be if distributed across the entire TKA population.

All analyses were performed using SAS v9.4 (Cary, NC). The probability of a type I error was set at .05, and all testing was 2 sided. Because patients cannot be identified, our institutional review board exempted this study from review.

Results

We identified 224,465 patients having TKA across all states participating in the NRD in 2014 (Table 1). The mean unadjusted 30-day TKA readmission rate was 4%. Several patient, clinical, payer, and hospital characteristics were associated with 30-day readmissions, including age, sex, median household income, comorbidities, index hospital length of stay, discharge destination, primary payer, location, hospital type, and hospital size (all $P < .01$).

The differences in readmission rates between Medicare and non-Medicare patients are demonstrated in Table 2. Although

Table 1
Demographic and Clinical Characteristics of Patients Undergoing Total Knee Arthroplasty.

Characteristics	Total Knee Arthroplasty		P Value ^a
	Not Readmitted	Readmitted ^e	
Number of patients (%)	216,649 (96)	7816 (4)	
Mean age (SD)	67 (9)	66 (10)	<.0001
Age group (%)			<.0001
45-54	22,338 (10)	662 (8)	
55-64	62,831 (29)	1791 (23)	
65-74	83,464 (38)	2748 (35)	
75-84	42,557 (20)	2178 (28)	
85 or older	5459 (3)	437 (6)	
Sex (%)			<.0001
Female	136,184 (63)	4456 (57)	
Male	80,465 (37)	3360 (43)	
Median household income, dollars (%)			<.0001
37,999 or less	43,462 (20)	1781 (23)	
38,000-47,999	56,382 (26)	2034 (27)	
48,000-63,999	58,611 (27)	2082 (27)	
64,000 or more	54,793 (23)	1787 (23)	
Comorbidities ^b (%)			
Diabetes without complications	41,850 (19)	1858 (24)	<.0001
Diabetes with complications	4651 (2)	269 (3)	<.0001
Chronic pulmonary disease	31,081 (14)	1507 (19)	<.0001
Rheumatologic disease	8778 (4)	376 (5)	.0009
Renal disease	10,199 (5)	760 (10)	<.0001
Congestive heart failure	4502 (2)	396 (5)	<.0001
Obese or overweight ^c	50,121 (23)	1882 (24)	.005
Length of stay (%)			<.0001
2 d or less	61,785 (28)	1581 (20)	
3 d	110,213 (51)	3810 (49)	
4 d	23,096 (11)	1101 (14)	
5 d or more	21,555 (10)	1324 (17)	
Discharge destination (%)			<.0001
Home	58,169 (27)	1515 (19)	
Home care	97,613 (45)	3079 (39)	
Skilled nursing facility	60,867 (28)	3222 (41)	
Primary payer (%):			<.0001
Medicare	127,495 (59)	5369 (69)	
Non-Medicare	89,154 (41)	2447 (31)	
Location (%)			<.0001
Urban	197,227 (91)	7093 (91)	
Rural ^d	19,422 (9)	723 (9)	
Hospital type (%)			<.0001
Teaching	122,897 (43)	3251 (42)	
Nonteaching	93,752 (57)	4565 (58)	
Hospital size (%)			<.0001
Small	38,395 (18)	1178 (15)	
Medium	58,406 (27)	2197 (28)	
Large	119,848 (55)	4441 (57)	

SD, standard deviation.

^a P values: t test for parametric continuous variables, Wilcoxon rank-sum test for nonparametric continuous variables, and chi-square test for proportions.

^b Represent 7 most common comorbidities.

^c Indicated by an International Classification of Diseases, Ninth Revision code.

^d Rural includes micropolitan areas.

^e Represents readmission within 30 d.

statistically significant, the 30-day readmission rates were clinically similar between the Medicare and non-Medicare patients (4% vs 3%). The majority of readmissions (57% of Medicare patients and 54% of non-Medicare patients) occurred after the first week from discharge. The majority of readmissions in both groups occurred with a length of stay of 3 days or more (80% for Medicare patients and 72% for non-Medicare patients, $P < .0001$). Medicare patients had a lower readmission rate on discharge at day 2 than non-Medicare patients (19% vs 28%, $P < .0001$). Medicare patients had a much higher rate of readmission from a skilled nursing facility than non-Medicare patients (48% vs 25%, $P < .0001$).

Adjusted predictors of readmission are demonstrated in Table 3. These results were based on a completed univariate analysis. In the

Table 2
Characteristics of Readmission Stratified by Payer.

Characteristics	Medicare	Non-Medicare	P Value ^a
Number of patients (%)	132,864 (59)	91,601 (41)	<.0001
Any readmission within 30 d (%)	5369 (4)	2447 (3)	<.0001
Time to readmission (d)			
Mean (SD)	11.5 (9)	11.2 (9)	.15
Median (IQR)	9 (4-18)	9 (4-18)	.11
Time to readmission group (%)			.07
1-7	2331 (43)	1119 (46)	
8-30	3038 (57)	1328 (54)	
Index length of stay, d (%)			<.0001
2 (or less)	842 (19)	567 (28)	
3	2693 (62)	1117 (56)	
4 (or more)	791 (18)	310 (16)	
Discharge destination (%)			<.0001
Home	844 (16)	671 (27)	
Home care	1924 (36)	1155 (47)	
Skilled nursing facility	2601 (48)	621 (25)	

IQR, interquartile range; SD, standard deviation.

^a P value: Wilcoxon rank-sum test for continuous variables and chi-square test for proportions.

adjusted analysis, male gender, decreased median household income, comorbidities, length of stay, Medicare payer, nonteaching hospital, rural hospital setting, and decreased hospital bed size were all associated with a higher likelihood of readmission (all $P < .05$).

The inpatient costs of readmission were examined and then extrapolated across the entire TKA population. The overall median cost for each readmission was \$6753 ± 175 (interquartile range). The mean cost of readmission was \$10,465 ± 15,257 (standard deviation). This was 36% of the overall total inpatient cost for the first 30 days from the index procedure. Extrapolating the readmission cost for the entire TKA population resulted in the readmission cost being 2% of the overall 30-day procedure cost.

Discussion

The Centers for Medicare and Medicaid Services' CJR bundle has focused efforts on improving quality of care for patients receiving a TKA. A major focus in assessing quality of care and reducing cost includes readmission rates. Our study has 3 main findings with implications for health policy related to arthroplasty readmissions. First, the overall readmission rate was low at 4%. Second, the greatest clinical predictors of readmission were renal disease, congestive heart failure, and length of stay greater than 4 days. Finally, readmissions comprised 36% of the actual cost of care in the initial 30 days for readmitted patients and 2% of the cost for all patients receiving a TKA.

In comparison to other orthopedic procedures, TKA readmission rates were low. This is in agreement with other national health service databases [13–21]. An analysis of the National Surgical Quality Improvement Program database demonstrated that the average 30-day readmission rates for arthroplasty, the highest volume procedure, was 2.8%. The readmission rate for TKAs is lower than that of other orthopedic procedures. For example, readmission rates for spine, posterior spinal instrumentation, and fusion for lumbar stenosis are approximately 10%, and rates for orthopedic procedures to address hip fractures are approximately 10%–15% [22].

In addition, TKA readmission rates are dramatically below rates for other common nonorthopedic surgical procedures. In cardiothoracic surgery, readmission rates following cardiac artery bypass grafting are 16% [23]. A range of colorectal procedures have readmissions between 10%–25% [24]. Bladder cancer patients undergoing a radical cystectomy have readmission rates around 25% [25].

Table 3
Multivariable Analysis of Each Predictor for 30-D Readmission vs No Readmission.

	Adjusted OR ^a (95% CI)	P Value
	Readmission vs No Readmission	
Age (y)		
45-54	Reference	
55-64	0.94 (0.86-1.03)	.20
65-74	0.90 (0.81-1.00)	.04
75-84	1.29 (1.16-1.44)	<.0001
85 or older	1.91 (1.66-2.20)	<.0001
Gender		
Female	Reference	
Male	1.36 (1.30-1.42)	<.0001
Household income		
\$37,999 or less	Reference	
\$38,000-\$47,999	0.90 (0.84-0.96)	.002
\$48,000-\$63,999	0.91 (0.85-0.97)	.004
\$64,000 or more	0.85 (0.80-0.91)	<.0001
Comorbidities		
Diabetes without complications	1.17 (1.10-1.23)	<.0001
Diabetes with complications	1.22 (1.07-1.40)	.003
Chronic pulmonary disease	1.35 (1.28-1.43)	<.0001
Rheumatologic disease	1.20 (1.07-1.33)	.001
Renal disease	1.55 (1.42-1.68)	<.0001
Congestive heart failure	1.59 (1.43-1.78)	<.0001
Obesity	1.07 (1.01-1.30)	.022
Length of stay (d)		
2 or less	Reference	
3	1.29 (1.21-1.37)	<.0001
4	1.69 (1.56-1.83)	.001
5 or more	1.98 (1.83-2.14)	<.0001
Medicare vs non-Medicare	1.27 (1.18-1.37)	<.0001
Teaching vs nonteaching	0.95 (0.90-0.99)	.03
Urban vs rural	1.06 (0.98-1.15)	.15
Hospital bed size		
Small	Reference	
Medium	1.20 (1.12-1.30)	<.0001
Large	1.15 (1.07-1.23)	<.0001

CI, confidence interval; OR, odds ratio.

^a Adjusted for patient and hospital characteristics (age, gender, median household income, comorbidities, index length of stay, primary payer, hospital type, and hospital size).

Other common procedures have readmission rates greater than TKA, including appendectomy at 6% [23]. The majority of these procedures are not elective as compared with TKA, resulting in an expected increase in readmissions as there are fewer options on patient selection for comorbidities. It would be expected that TKA has lower readmission rates. Here, these other procedures give an interesting perspective in the unique challenges required to lower readmission rates for TKA, where the rate of readmission is already very low.

Other groups have tried to identify modifiable and non-modifiable predictors of readmission to identify patients at increased risk, determine strategies to decrease risk, and set patient expectations in arthroplasty procedures. In our study, the greatest predictors of readmission were renal disease, congestive heart failure, and length of stay greater than 4 days (all had odds ratios >1.5). These modifiable medical comorbidities suggest the importance of preoperative optimization to lower readmission rates. Patients with an increased length of stay should be identified as a high-risk group and followed closely after discharge. Many other large, national databases and institutional studies have observed the strong predictor in length of stay on readmission rates [26–29]. A longer length of stay implies that a patient had a more complicated course during their admission [30].

The interest in lowering TKA readmission rates is based on the logic that decreased readmissions provide increased quality of care. This assumes that, as readmissions decrease, patient outcomes

improve and costs of each episode of care decrease. In our study, we analyzed the cost of care associated with each readmission. The mean cost of each readmission per readmitted patient was \$10,465 and is in agreement with other data that report a TKA readmission cost of \$10,200 [23]. In the readmitted population, this is one-third of the overall cost; a major contributor of the final cost of care. Given these factors, distributing the cost of readmission across the entire population of individuals receiving a TKA results in a cost of readmission per all patients receiving a TKA of \$365, only 2% of overall cost. This suggests from a cost perspective that spending more than this limit per patient to reduce readmission rates results in less cost savings. Our cost-benefit analysis suggests that this limit is being approached. Previous work is in alignment with these findings. Based on institutional charge data, other groups have determined that the cost burden of TKA is approximately 2.8%. This suggests that the profit margin would have to be greater than 2.8% for the hospital to recover the cost of the procedure and all readmissions related to the procedure.

Similar to all research based on large data sets, this study is dependent on the accuracy of coding and recorded patient data. Our findings should be interpreted in the context of these limitations. We are examining readmission using administrative data, which does not contain all factors that could impact readmission, such as family support or driving distance from the hospital. However, we adjust for several factors that can influence readmissions, such as comorbidities. In addition, by using the NRD, we are able to examine readmissions for TKA at a national level across all payers. To date, most studies examining TKA readmissions are based on Medicare databases or other administrative databases restricted to a single state or group of states. Our ability to conduct a complete cost analysis is limited as outpatient charge data after inpatient care was unavailable. Follow-up care has been reported to be approximately one-third of the total cost of TKA [31]. Finally, our objective was to look at 30-day readmission rates. If analysis was extended to 90 days, the cost of readmissions would increase. Even with a conservative estimate of doubling readmission rates and cost, readmission cost would still be approximately 4% of the overall cost of the TKA for the entire population.

Hospital readmissions are being used as a key performance indicator to gauge the quality of care provided and to help control cost. Some hospital readmissions are avoidable, suggesting sub-optimal initial care, but the validity of this assumption has been questioned [32–34]. Our study provides evidence that questions the focus on readmissions in this population. The overall low readmission rate as compared with other targeted diagnoses provides a limited return in quality improvement. Further from a cost perspective, readmission costs are a small portion of the overall cost (2%), given this low readmission rate. This implies that as the limit in these areas are being reached, other drivers of quality and cost that have larger rooms for improvement should be considered. Minimizing readmission rates are important for quality. Identifying the main factors for the other 98% of cost will result in significant improvement in quality and cost.

References

- [1] Services CfMM. Bundled payments for care improvement (BPCI) initiative.
- [2] Burwell SM. Setting value-based payment goals—HHS efforts to improve U.S. health care. *N Engl J Med* 2015;372:897–9.
- [3] Saucedo JM, Marecek GS, Wanke TR, Lee J, Stulberg SD, Puri L. Understanding readmission after primary total hip and knee arthroplasty: who's at risk? *J Arthroplasty* 2014;29:256–60.
- [4] Froimson MI, Rana A, White Jr RE, Marshall A, Schutzer SF, Healy WL, et al. Bundled payments for care improvement initiative: the next evolution of payment formulations: AAHKS bundled payment task force. *J Arthroplasty* 2013;28(8 Suppl):157–65.

- [5] Dundon JM, Bosco J, Slover J, Yu S, Sayeed Y, Iorio R. Improvement in total joint replacement quality metrics: year one versus year three of the bundled payments for care improvement initiative. *J Bone Joint Surg Am* 2016;98:1949–53.
- [6] D'Apuzzo M, Westrich G, Hidaka C, Jung Pan T, Lyman S. All-cause versus complication-specific readmission following total knee arthroplasty. *J Bone Joint Surg Am* 2017;99:1093–103.
- [7] Healthcare Cost and Utilization Project (HCUP). NRD Database Overview. <https://www.hcup-us.ahrq.gov/nrdoverview.jsp> [accessed 27.04.18].
- [8] Mesko NW, Bachmann KR, Kovacevic D, LoGrasso ME, O'Rourke C, Froimson MI. Thirty-day readmission following total hip and knee arthroplasty - a preliminary single institution predictive model. *J Arthroplasty* 2014;29:1532–8.
- [9] Deyo RA, Cherkin DC, Ciol MA. Adapting a clinical comorbidity index for use with ICD-9-CM administrative databases. *J Clin Epidemiol* 1992;45:613–9.
- [10] Trinh QD, Bianchi M, Sun M, Sammon J, Schmitges J, Shariat SF, et al. Discharge patterns after radical prostatectomy in the United States of America. *Urol Oncol* 2013;31:1022–32.
- [11] Yu HY, Hevelone ND, Lipsitz SR, Kowalczyk KJ, Hu JC. Use, costs and comparative effectiveness of robotic assisted, laparoscopic and open urological surgery. *J Urol* 2012;187:1392–9.
- [12] Sachs T, Pomposelli F, Hamdan A, Wyers M, Schermerhorn M. Trends in the national outcomes and costs for claudication and limb threatening ischemia: angioplasty vs bypass graft. *J Vasc Surg* 2011;54:1021–1031.e1.
- [13] Adelani MA, Keeney JA, Nunley RM, Clohisy JC, Barrack RL. Readmission following total knee arthroplasty: venous thromboembolism as a “never event” is a counterproductive misnomer. *J Arthroplasty* 2013;28:747–50.
- [14] Avram V, Petrucelli D, Winemaker M, de Beer J. Total joint arthroplasty readmission rates and reasons for 30-day hospital readmission. *J Arthroplasty* 2014;29:465–8.
- [15] Huddleston JI, Maloney WJ, Wang Y, Verzier N, Hunt DR, Herndon JH. Adverse events after total knee arthroplasty: a national medicare study. *J Arthroplasty* 2009;24(6 Suppl):95–100.
- [16] Schaeffer JF, Scott DJ, Godin JA, Attarian DE, Wellman SS, Mather 3rd RC. The association of ASA class on total knee and total hip arthroplasty readmission rates in an academic hospital. *J Arthroplasty* 2015;30:723–7.
- [17] Tayne S, Merrill CA, Smith EL, Mackey WC. Predictive risk factors for 30-day readmissions following primary total joint arthroplasty and modification of patient management. *J Arthroplasty* 2014;29:1938–42.
- [18] Zmistowski B, Restrepo C, Hess J, Adibi D, Cangoz S, Parvizi J. Unplanned readmission after total joint arthroplasty: rates, reasons, and risk factors. *J Bone Joint Surg Am* 2013;95:1869–76.
- [19] Ravi B, Croxford R, Austin PC, Hollands S, Paterson JM, Bogoch E, et al. Increased surgeon experience with rheumatoid arthritis reduces the risk of complications following total joint arthroplasty. *Arthritis Rheum* 2014;66:488–96.
- [20] Singh JA, Lu X, Rosenthal GE, Ibrahim S, Cram P. Racial disparities in knee and hip total joint arthroplasty: an 18-year analysis of national Medicare data. *Ann Rheum Dis* 2014;73:2107–15.
- [21] Ramkumar PN, Chu CT, Harris JD, Athiviraham A, Harrington MA, White DL, et al. Causes and rates of unplanned readmissions after elective primary total joint arthroplasty: a systematic review and meta-analysis. *Am J Orthop (Belle Mead NJ)* 2015;44:397–405.
- [22] Ali AM, Gibbons CE. Predictors of 30-day hospital readmission after hip fracture: a systematic review. *Injury* 2017;48:243–52.
- [23] Qasim M, Andrews RM. Post-surgical readmissions among patients living in the poorest communities, 2009: Statistical Brief #142. In: *Healthcare cost and utilization project (HCUP) Statistical Briefs*. Rockville (MD); 2006.
- [24] Damle RN, Alavi K. Risk factors for 30-d readmission after colorectal surgery: a systematic review. *J Surg Res* 2016;200:200–7.
- [25] Hu M, Jacobs BL, Montgomery JS, He C, Ye J, Zhang Y, et al. Sharpening the focus on causes and timing of readmission after radical cystectomy for bladder cancer. *Cancer* 2014;120:1409–16.
- [26] Paxton EW, Inacio MC, Singh JA, Love R, Bini SA, Namba RS. Are there modifiable risk factors for hospital readmission after total hip arthroplasty in a US healthcare system? *Clin Orthop Relat Res* 2015;473:3446–55.
- [27] Valensi P, Perret G, Vassy R, Uzzan B, Nicolas P, Attali JR. Effect of nifedipine on thyrotropin, prolactin, and thyroid hormone release in man: a placebo-controlled study. *Fundam Clin Pharmacol* 1989;3:59–66.
- [28] Mesko NW, Bachmann KR, Kovacevic D, LoGrasso ME, O'Rourke C, Froimson MI. Thirty-day readmission following total hip and knee arthroplasty - a preliminary single institution predictive model. *J Arthroplasty* 2014;29:1532–8.
- [29] Kurtz SM, Lau EC, Ong KL, Adler EM, Kolisek FR, Manley MT. Which hospital and clinical factors drive 30- and 90-day readmission after TKA? *J Arthroplasty* 2016;31:2099–107.
- [30] El Bitar YF, Illingworth KD, Scaife SL, Horberg JV, Saleh KJ. Hospital length of stay following primary total knee arthroplasty: data from the Nationwide inpatient sample database. *J Arthroplasty* 2015;30:1710–5.
- [31] Nichols CI, Vose JG. Clinical outcomes and costs within 90 Days of primary or revision total joint arthroplasty. *J Arthroplasty* 2016;31:1400–1406.e3.
- [32] Weissman JS, Ayanian JZ, Chasan-Taber S, Sherwood MJ, Roth C, Epstein AM. Hospital readmissions and quality of care. *Med Care* 1999;37:490–501.
- [33] Jimenez-Puente A, Garcia-Alegria J, Gomez-Aracena J, Hidalgo-Rojas L, Lorenzo-Nogueiras L, Perea-Milla-Lopez E, et al. Readmission rate as an indicator of hospital performance: the case of Spain. *Int J Technol Assess Health Care* 2004;20:385–91.
- [34] Remes V. CORR insights ((R)): are there modifiable risk factors for hospital readmission after total hip arthroplasty in a US healthcare system? *Clin Orthop Relat Res* 2015;473:3456–7.